



QUAT-STAT™ SC



One-Step Disinfectant • Cleaner

A heavy duty disinfectant cleaner that cleans, disinfects, and deodorizes in one labor saving step. A concentrated Hospital Use disinfectant cleaner that is effective against a broad spectrum of bacteria, is *virucidal, fungicidal, and inhibits the growth of mold and mildew and their odors, when used as directed.



Uses

- Floors and walls
- Formica® and Fibreglas®
- Doorknobs
- Toilets and urinals
- Metal
- Porcelain
- Finished woodwork
- Desks, counter tops and chairs
- Telephones
- Shower stalls, tubs and tiles
- Glazed porcelain and glazed ceramic
- Plastic
- Stainless steel enamel

Benefits

- One step cleaner disinfectant
- Kills HIV-1 (AIDS Virus), HBV (Hepatitis B Virus), VRE (Vancomycin Resistant Enterococcus faecalis), MRSA (Methicillin Resistant Staphylococcus Aureus).

Properties

Color	Purple
Detergency	Excellent
Film residue	None
Flammability	Nonflammable
Fragrance	Fresh
pH	12.0 - 13.5
Shelf life (of concentrated product).....	1 year
Octyl decyl dimethyl ammonium chloride	6.510%
Diocetyl dimethyl ammonium chloride.....	2.604%
Didecyl dimethyl ammonium chloride.....	3.906%
Alkyl (C ₁₄ , 50%; C ₁₂ , 40%; C ₁₆ , 10%) dimethyl benzyl ammonium chloride	8.680%

Safety

KEEP OUT OF REACH OF CHILDREN. **DANGER:** FOR HOSPITAL, INSTITUTIONAL AND INDUSTRIAL USE. Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin, or on clothing. Wear protective eyewear (goggles, face shield or safety glasses), protective clothing and protective (rubber or chemical resistant gloves). Harmful if swallowed and/or if absorbed through the skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing before reuse. **FIRST AID:** IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing eye. IF ON SKIN OR CLOTHING: Take off contaminated clothing and wash clothing before reuse. Rinse skin immediately with plenty of water for 15 - 20 minutes. IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor or going for treatment.

Available In

FASTDRAW™ SYSTEM Case/4 - 2 L Bottles: Item #34047-00
 GREEN EARTH® II SYSTEM Case/2 - 4 L Boxes: Item #34037-00
 FASTDOSE™ SYSTEM Case/6 Quarts (946 mL) Bottles: Item #34048-00

Acceptance

Registered with the U.S. Environmental Protection Agency. EPA Reg. No. 6836-78-4170.

Directions For Use

Read the entire label before using this product.

For Hospitals, Medical and Dental Offices, Nursing Homes and other Medical Facilities: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **Dilution:** Disinfection 1:256 ½ oz. per gallon of water.

DISINFECTION/VIRUCIDAL*/FUNGICIDAL/MOLD AND MILDEW CONTROL. Apply use-solution to hard, nonporous surfaces, thoroughly wetting surfaces with a cloth, mop, sponge or sprayer or by immersion. Treated surfaces must remain wet for 10 minutes. Wipe dry with a cloth, sponge, or mop or allow to air dry. For heavily soiled areas, a preliminary cleaning is required. For sprayer applications, use a coarse spray device. Spray 6 - 8 inches from the surface; rub with a brush, sponge or cloth. Do not breathe spray. Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected. Rinse all surfaces that come in contact with food such as countertops, exteriors of appliances, tables and stovetops with potable water before reuse. Do not use on utensils, glassware and dishes. For fungicidal activity, this product is an effective fungicide against Trichophyton Mentagrophytes [the athlete's foot fungus] in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities. This product is a one-step fungicide at ½ oz. per gallon in 250 ppm hard water and 5% serum. At ½ oz. per gallon in 400 ppm hard water, this product must be used on pre-cleaned surfaces. For mold and mildew, at ½ oz. per gallon, this product will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, nonporous surfaces. Follow disinfection instructions. Repeat treatment every seven days, or more often if new growth appears. **BACTERICIDAL STABILITY OF USE-DILUTION:** The use solution of this product, when stored in a sealed container such as a spray bottle, remains effective for up to 1 year. If product becomes visibly dirty or contaminated, the use-dilution must be discarded and fresh product prepared. Always use clean, properly labeled containers when diluting this product. Bactericidal stability of the use-dilution does not apply to open containers such as buckets or pails. This product is not for use on medical device surfaces. **NON-ACID TOILET BOWL AND URINAL DISINFECTANT CLEANER DIRECTIONS:** Remove gross filth prior to disinfection. From Concentrate: Add ½ oz. to the bowl and mix. Brush over exposed surfaces and under the rim, allow to stand for 10 minutes and flush. From Use Solution: Empty toilet bowl or urinal and apply ½ oz. per gallon use-solution to exposed surfaces and under the rim with a cloth, mop, sponge or mechanical spray. Brush or swab thoroughly, allow to stand for 10 minutes and flush. For Heavy Duty Use: Empty toilet bowl or urinal and apply 2 oz. per gallon use-solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer, brush or swab thoroughly, allow to stand for 10 minutes and flush. ***KILLS HIV AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV). **SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OR HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.** **Personal protection:** Clean-up must always be done wearing protective latex gloves, gowns, masks and eye protection. **Cleaning Procedure:** Blood and other body fluids containing HIV or HBV or HCV must be thoroughly cleaned from surfaces and objects before application of this product. **Contact Time:** Leave surface wet for 1 minute (60 seconds) for HIV-1 and 10 minutes for HBV and HCV. Use a 10-minute contact time for disinfection against all other viruses, bacteria and fungi claimed. **Disposal of infectious material:** Blood, body fluids, cleaning materials and clothing should be autoclaved and disposed of according to local regulations for infectious waste disposal.

Literature Item #34092-92

